

SYMPLECTIC 4-MANIFOLDS WITH ARBITRARY FUNDAMENTAL GROUP NEAR THE BOGOMOLOV-MIYAOKA-YAU LINE

SCOTT BALDRIDGE AND PAUL KIRK

ABSTRACT. In this paper we construct a family of symplectic 4-manifolds with positive signature for any given fundamental group G that approaches the BMY line. The family is used to show that one cannot hope to do better than the BMY inequality in finding a lower bound for the function $f = \chi + b\sigma$ on the class of all minimal symplectic 4-manifolds with a given fundamental group.

1. INTRODUCTION

Let $\chi(S)$ and $\sigma(S)$ denote the Euler characteristic and signature of a 4-manifold respectively. Minimal complex surfaces S of general type satisfy $c_1^2(S) > 0$, $\chi(S) > 0$ and

$$2\chi_h(S) - 6 \leq c_1^2(S) \leq 9\chi_h(S)$$

where $c_1^2(S) = 2\chi(S) + 3\sigma(S)$ and $\chi_h(S) = \frac{1}{4}(\chi(S) + \sigma(S))$. The second inequality is usually referred to as the Bogomolov-Miyaoka-Yau inequality. Finding symplectic (or Kähler) 4-manifolds on or near the BMY line has a long and interesting history (c.f. [8], [2], [3], [9], [5], [10], [11]). In particular this means looking for symplectic 4-manifolds with positive signature.

All known examples of symplectic 4-manifolds on the BMY line have large fundamental groups. In fact, if S is a complex surface differing from $\mathbb{CP}^2$ (the only known simply-connected example on the BMY line), the equality $c_1^2(S) = 9\chi_h(S)$ holds if and only if the unit disk $D^4 = \{(z_1, z_2) \in \mathbb{C}^2 \mid |z_1|^2 + |z_2|^2 \leq 1\}$ covers S [6], [13], [7]. This implies for such S that $|\pi_1(S)| = \infty$. The goal has generally been to produce examples that fill in the geography with respect to (c_1^2, χ_h) . In this paper we are interested what can be said for a given fundamental group.

Stipsicz constructed simply connected symplectic 4-manifolds C_n for which $c_1^2(C_n)/\chi_h(C_n) \rightarrow 9$ as $n \rightarrow \infty$ in [11]. Our main theorem generalizes this result to any fundamental group.

Theorem 1. *Let G have a presentation with g generators $x_1, \dots, x_g$ and r relations $w_1, \dots, w_r$. For each integer $n > 1$, there exists a symplectic*

The first author gratefully acknowledges support from the NSF grant DMS-0507857. The second author gratefully acknowledges support from the NSF grant DMS-0202148.

4-manifold $M(G, n)$ with fundamental group G with Euler characteristic

$$\chi(M(G, n)) = 75n^2 + 256n + 130 + 12(g + r + 1),$$

and signature

$$\sigma(M(G, n)) = 25n^2 - 68n - 78 - 8(g + r + 1).$$

Our interest in this question developed while investigating pairs $(a, b) \in \mathbb{R}^2$ for which the function $f = a\chi + b\sigma$ has a lower bound on the class of symplectic manifolds with a given fundamental group (see [1]). In that article we considered the following.

Fix a finitely presented group G and let $\mathfrak{M}$ denote either the class $\mathfrak{M}(G)$ of closed symplectic 4-manifolds with fundamental group G or the class $\mathfrak{M}^{\min}(G)$ of minimal, closed symplectic 4-manifolds with fundamental group G .

For $b \in \mathbb{R}$, define $f_{\mathfrak{M}}(b) \in \mathbb{R} \cup \{-\infty\}$ to be the infimum

$$f_{\mathfrak{M}}(b) = \inf_{M \in \mathfrak{M}} \{\chi(M) + b\sigma(M)\}.$$

(In [1] we considered the infimum $f_{\mathfrak{M}}(a, b)$ of $a\chi + b\sigma$ on $\mathfrak{M}$ and showed that if $a \leq 0$ the infimum is $-\infty$. Thus we restrict to $f_{\mathfrak{M}}(1, b)$, which we more compactly denote by $f_{\mathfrak{M}}(b)$ in the present article.)

We showed in [1] that the set

$$D_{\mathfrak{M}} = \{b \mid f_{\mathfrak{M}}(b) \neq -\infty\}$$

(the *domain* of $f_{\mathfrak{M}}$) is an interval satisfying

$$[-1, 1] \subset D_{\mathfrak{M}(G)} \subset (-\infty, 1] \text{ and } [-1, \frac{3}{2}] \subset D_{\mathfrak{M}^{\min}(G)} \subset (-\infty, \frac{3}{2}].$$

The upper bounds are sharp; in fact $1 \in D_{\mathfrak{M}(G)}$ and $\frac{3}{2} \in D_{\mathfrak{M}^{\min}(G)}$.

We are interested in the value of the left endpoint e_G of $D_{\mathfrak{M}(G)}$, which is an intriguing invariant of a group G . (It may or may not be contained in $D_{\mathfrak{M}(G)}$.) Since $e_G \leq -1$, a straightforward argument shows that e_G is also the left endpoint of $D_{\mathfrak{M}^{\min}(G)}$.

In [1] we observed that the results of Stipcitz gives a better lower bound (than $-\infty$) when G is the trivial group, and so a consequence of the result of this article is an extension to all G . In fact Theorem 1 easily implies the following corollary.

Corollary 2. *For any finitely presented group G ,*

$$D_{\mathfrak{M}(G)} \subset [-3, 1] \text{ and } D_{\mathfrak{M}^{\min}(G)} \subset [-3, \frac{3}{2}].$$

□

The BMY inequality $c_1^2 \leq 9\chi_h$ is equivalent to $f_{\mathfrak{M}^{\min}(G)}(-3) \geq 0$ provided G is not a surface group. Hence the BMY conjecture and Corollary 2 together imply that $e_G = 3$. Thus, a weaker form of the BMY conjecture could be stated as follows.

Conjecture 3 (Weak BMY Conjecture). *For each finitely presented group G , $e_G = -3$.*

2. THE CONSTRUCTION

We use the following notation. If X and Y are symplectic 4-manifolds containing symplectic surfaces $F_X \subset X$ and $F_Y \subset Y$ such that $F_X^2 + F_Y^2 = 0$, then the symplectic sum ([4]) of X and Y along F_X and F_Y will be denoted by

$$X \#_{F_X, F_Y} Y.$$

Recall that topologically $X \#_{F_X, F_Y} Y$ is obtained by removing tubular neighborhoods of F_X and F_Y and identifying the resulting boundaries in a fiber-preserving way. Moreover, if $E_X \subset X$ (resp. $E_Y \subset Y$) is a symplectic surface intersecting F_X once transversally (resp. intersecting F_Y transversally), then the fiber sum can be constructed so that (the connected sum) $E_X \# E_Y$ is a symplectic surface in $X \#_{F_X, F_Y} Y$.

2.1. The first piece: symplectic manifolds with given fundamental group. The following theorem was proven in [1].

Theorem 4. *Let G have a presentation with g generators $x_1, \dots, x_g$ and r relations $w_1, \dots, w_r$. Then there exists a minimal symplectic 4-manifold $M(G)$ with $\pi_1(M(G)) \cong G$, Euler characteristic $\chi(M(G)) = 12(g + r + 1)$, and signature $\sigma(M(G)) = -8(g + r + 1)$.*

We will use the following observation. The manifold $M(G)$ constructed in Theorem 4 is obtained by taking fiber sums of a certain base manifold with $g + r + 1$ copies of the basic elliptic surface $E(1)$. Since $E(1)$ admits a singular fibration with symplectic generic fibers and 6 cusp fibers (which are simply connected), so does $E(1) - F$, where F denotes the generic fiber in $E(1)$ along which the fiber sum giving M is constructed. Thus each $M(G)$ contains a symplectic torus T_0 such that the induced homomorphism $\pi_1(T_0) \rightarrow \pi_1(M(G))$ is trivial.

2.2. The second piece: symplectic manifolds near the BMY line. In [11], Stipsicz proved the following theorem.

Proposition 5 (Stipsicz). *For each non-negative integer n , there exists a symplectic 4-manifold $X(n)$ which admits a genus- $(15n+1)$ Lefschetz fibration with a section T_{n+2} of genus $(n+2)$ and self-intersection $-(n+1)$. Furthermore, $X(n)$ can be equipped with a symplectic structure such that T_{n+2} is a symplectic submanifold. The projection map $X(n) \rightarrow T_{n+2}$ induces an isomorphism on fundamental groups. The Euler characteristic of $X(n)$ is $\chi(X(n)) = 75n^2 + 180n + 12$ and the signature is $\sigma(X(n)) = 25n^2 - 60n - 8$.*

Denote by $F_{15n+1} \subset X(n)$ a fixed generic fiber of $X(n)$. This is a symplectic surface with trivial normal bundle.

2.3. The third piece: a simply connected manifold. Gompf constructs a symplectic 4-manifold $S_{1,1}$ in [4, Lemma 5.5] which contains a disjoint pair T, F of symplectically embedded surfaces T of genus one and F of genus two,

with trivial normal bundles such that $S_{1,1} - (T \cup F)$ is simply connected. Thus the symplectic sum A of two copies $S_{1,1}$ along the genus two surfaces

$$A = S_{1,1} \#_{F,F} S_{1,1}$$

contains a pair of disjointly embedded symplectic tori $T_1 \cup T_2 \subset A$ with trivial normal bundles so that the complement $A - (T_1 \cup T_2)$ is simply connected. Since $S_{1,1}$ has Euler characteristic 23 and signature -15 , $\chi(A) = 50$ and $\sigma(A) = -30$.

The manifold A has a useful property, whose proof is a simple application of the Seifert-Van Kampen theorem.

Proposition 6. *Suppose B and C are symplectic 4-manifolds containing symplectic tori $i_B : T_B \subset B$ and $i_C : T_C \subset C$ with trivial normal bundles.*

Let $D = B \#_{T_B, T_1} A \#_{T_2, T_C} C$ be the fiber sum of B , A , and C . Then

$$\pi_1(D) = (\pi_1(B)/N((i_B)_*(\pi_1(T_B)))) \star (\pi_1(C)/N((i_C)_*(\pi_1(T_C))))$$

where $\star$ denotes free product and $N(H)$ denotes the normal closure of a subgroup H . $\square$

2.4. The fourth piece: “Elbows”. Let T be a torus and $\{a, b\}$ a pair of smoothly embedded loops forming a symplectic basis of $\pi_1 T$. Let $\varphi : T \rightarrow T$ be the Dehn twist around a . The mapping torus Y_φ fibers over S^1 with fiber T . Let $t_1 : S^1 \rightarrow Y_\varphi$ denote a section. Taking a product of Y_φ with S^1 yields a symplectic 4-manifold $Y_\varphi \times S^1$ (this is just Thurston’s manifold from [12]) which fibers over a torus with symplectic torus fibers. Moreover, the symplectic structure can be chosen so that the section $t_1 \times \text{id} : S^1 \times S^1 \rightarrow Y_\varphi \times S^1$ is symplectic. Denote by $s_1 : S^1 \rightarrow \{p\} \times S^1 \subset Y_\varphi \times S^1$ the loop representing the second factor.

Note that $Y_\varphi \times S^1$ contains a torus $T' = b \times s_1$, where b is the curve described above in the fiber of Y_φ . The torus T' is homologically non-trivial by the Kunneth theorem, since b is non-trivial in $H_1(Y_\varphi)$, and is Lagrangian with respect to the symplectic structure on $Y_\varphi \times S^1$. Thus the symplectic structure on $Y_\varphi \times S^1$ can be perturbed slightly to make T' symplectic. Note moreover that T' is disjoint from the section $t_1 \times s_1 : S^1 \times S^1 \rightarrow Y_\varphi \times S^1$ since we can assume that t_1 intersects the fiber containing b in a point which does not lie on b . The tubular neighborhood of T' in $Y_\varphi \times S^1$ is trivial since b can be isotoped off itself in a fiber of $Y_\varphi \rightarrow S^1$. Similarly the tubular neighborhood of the section $t_1 \times s_1$ is trivial since t_1 can be pushed off itself in Y_φ .

Define $Elb(n)$ to be the fiber sum $Elb(n) = (Y_\varphi \times S^1) \#_{T, T^2} (T^2 \times \Sigma_{n-1})$. The fiber sum can be carried out so that the sections of $Y_\varphi \times S^1 \rightarrow S^1 \times S^1$ and $T^2 \times \Sigma_{n-1} \rightarrow \Sigma_{n-1}$ yield a symplectic section of the resulting fibration $Elb(n) \rightarrow \Sigma_n$. Thus $Elb(n)$ contains a disjoint pair of symplectic surfaces with trivial normal bundles, a torus $T' = b \times s_1$ and a genus n surface, the image of the section, which we denote by D_n .

Letting $t_2, s_2, \dots, t_n, s_n$ denote the generators of $\pi_1(\Sigma_{n-1})$, one computes

$$\pi_1(Elb(n)) = \langle a, b, t_1, s_1, \dots, t_n, s_n \mid \begin{array}{l} a \text{ central}, [b, t_1] = a, \\ [b, t_i] = 1 \text{ for } i > 1, [b, s_i] = 1 \text{ for all } i, \\ \prod_{i=1}^n [t_i, s_i] = 1 \end{array} \rangle.$$

The inclusion of T' into $Elb(n)$ takes the generators of $\pi_1 T'$ to b and s_1 , and the inclusion of D_n takes the standard basis to $t_1, s_1, \dots, t_n, s_n$. The Euler characteristic and signature of $Elb(n)$ both vanish.

The manifold $Elb(n) - D_n$ is a punctured torus fibration over Σ_n , and hence has a presentation with the same generators and all the same relations except that one no longer has a commuting with b . i.e. a commutes with all generators except b .

2.5. The fifth piece: an elliptic surface. We find a symplectically embedded surface J of genus $n + 3$ and self-intersection $n + 1$ in the elliptic surface $E(n + 5)$ such that $E(n + 5) - J$ is simply connected as follows. Consider $n + 3$ copies of the generic fiber and one copy of the section in a fibration $E(n + 5) \rightarrow \mathbb{CP}^1$ with $6(n + 5)$ cusp fibers. The section and fibers are symplectic with regards to the symplectic structure on the elliptic fibration $E(n + 5)$. Resolve the $n + 3$ transverse double points ([4]) to get a symplectically embedded surface J of genus $n + 3$ and self-intersection $n + 1$ (the fiber hits the section once and that section has self-intersection $-(n + 5)$). The complement $E(n + 5) - J$ is simply connected because $E(n + 5)$ has a simply connected fiber which intersects J in one point: the normal circle of a tubular neighborhood of J is nullhomotopic in $E(n + 5) - J$.

2.6. Putting the pieces together. We begin by a modification of Stipicz's construction. Let $Z(n)$ be the fiber sum of $Elb(15n + 1)$ and $X(n)$ along $D_{15n+1} \subset Elb(15n + 1)$ and the fiber F_{15n+1} of the Lefschetz fibration $X(n) \rightarrow \Sigma_{n+2}$

$$Z(n) = Elb(15n + 1) \#_{D_{15n+1}, F_{15n+1}} X(n).$$

The fiber sum can be constructed so that the fiber $T \subset Elb(15n + 1)$ and the section $T_{n+2} \subset X(n)$ add to yield a symplectic surface of genus $n + 3$, $K_{n+3} = T \# T_{n+2} \subset Z(n)$ [4]. The important property of $Z(n)$ is that it contains a symplectic torus T' , since D_{15n+1} and T' are disjoint.

The fundamental group of $Z(n)$ is easily computed, since $Elb(15n + 1) - D_{15n+1}$ is a fiber bundle with punctured torus fibers and $X(n) - F_{15n+1}$ is a Lefschetz fibration over a punctured genus $n + 2$ surface with at least one simply connected fiber. Using the Seifert-Van Kampen theorem and Novikov additivity one obtains the following.

Lemma 7. *The fundamental group of $Z(n)$ is the free product of $\mathbb{Z}$ with generator b , and a genus $n + 2$ surface group generated by x_i, y_i :*

$$\pi_1(Z(n)) = \mathbb{Z}b \star \langle x_i, y_i, i = 1, \dots, n + 2 \mid \prod [x_i, y_i] = 1 \rangle.$$

Moreover, $Z(n)$ contains a disjoint pair of symplectic surfaces, $T' \cup K_{n+3} \subset Z(n)$ satisfying $[T']^2 = 0$, and $[K_{n+3}]^2 = -n - 1$. The induced homomorphism $\pi_1(T') \rightarrow \pi_1(Z(n))$ is the map

$$\langle a, s_1 \mid [a, s_1] \rangle \rightarrow \pi_1 Z(n) \quad a \mapsto a, s_1 \mapsto 1.$$

The induced homomorphism $\pi_1(K_{n+3}) \rightarrow \pi_1(Z(n))$ is the map

$$\langle a, b, x_1, y_1, \dots, x_{n+2}, y_{n+2} \mid [a, b] \prod [x_i, y_i] = 1 \rangle \rightarrow \pi_1(Z(n))$$

$$a \mapsto a, b \mapsto 1, x_i \mapsto x_i, y_i \mapsto y_i.$$

Moreover, $\chi(Z(n)) = 75n^2 + 240n + 12$ and $\sigma(Z(n)) = 25n^2 - 60n - 8$. $\square$

The symplectic sum of $Z(n)$ with $E(n+5)$ along J , $Z(n) \#_{K_{n+3}, J} E(n+5)$ is a simply connected symplectic 4-manifold containing a torus T_1 with trivial normal bundle and appropriate Euler characteristic and signature. We take symplectic sum of this manifold with A to obtain an example with a torus whose complement is simply connected.

Define $W(n)$ to be the symplectic sum

$$W(n) = A \#_{T_1, T'} Z(n) \#_{K_{n+3}, J} E(n+5).$$

Then since $\pi_1(A - (T_2 \cup T_2)) = 1$, the following proposition follows straightforwardly.

Proposition 8. *The symplectic manifold $W(n)$ is simply connected and contains a symplectic torus $T_2 \subset W(n)$ with trivial normal bundle so that $\pi_1(W(n) - T_2) = 1$. It has Euler characteristic $\chi(W(n)) = 75n^2 + 256n + 130$ and signature $\sigma(W(n)) = 25n^2 - 68n - 78$. $\square$*

We can now prove Theorem 1.

Proof of Theorem 1. The symplectic sum

$$M(G, n) = M(G) \#_{T_0, T_2} W(n)$$

has fundamental group G by Proposition 6. The calculations of $\chi(M(G, n))$ and $\sigma(M(G, n))$ are routine. $\square$

REFERENCES

- [1] S. Baldridge and P. Kirk, *On symplectic 4-manifolds with prescribed fundamental group*, May 2005, E-print GT0504345.
- [2] Z. Chen, *On the geography of surfaces — simply connected minimal surfaces with positive index*, Math. Ann. **126** (1987), 141-164.
- [3] Z. Chen, *The existence of algebraic surfaces with preassigned Chern numbers*, Math. Z. **206** (1991), 241-254.
- [4] R. Gompf, *A new construction of symplectic manifolds*. Ann. of Math. **142** (1995), 527-595.
- [5] R. Gompf and A. Stipsicz, '4-Manifolds and Kirby Calculus.' Graduate Studies in Mathematics **20**, American Mathematical Society, Providence, Rhode Island, 1999.
- [6] F. Hirzebruch, *The signature of ramified coverings*, in Global Analysis (Papers in Honor of K. Kodaira), (D. Spencer and S. Iyanaga, eds.), Princeton University Press 1969, 253-265.

- [7] Y. Miyaoka, *The minimal number of quotient singularities on surfaces with given numerical invariants*. Math. Ann. **268** (1984), 159–171.
- [8] B. Moishezon and M. Teicher, *Simply connected algebraic surfaces of positive index*. Invent. Math. **89** (1987), 601–644.
- [9] U. Persson, C. Peters, and G. Xiao, *Geography of spin surfaces*. Topology **35** (1996), 845–862.
- [10] A. Stipsicz, *Simply connected 4-manifolds near the Bogomolov-Miyaoka-Yau line*. Math. Res. Lett. **5** (1998), 723–730.
- [11] A. Stipsicz, *Simply connected symplectic 4-manifolds with positive signature*, Tr. J. Mathematics, Proceedings of 6th Gökova Geometry-Topology Conference, **23** (1999), 145–150.
- [12] W. P. Thurston, *Some simple examples of symplectic manifolds*. Proc. Amer. Math. Soc. **55** (1976), no. 2, 467–468.
- [13] S.-T. Yau, *Calabi’s conjecture and some new results in algebraic geometry* Proc. nat. Acad. Sci. USA **74** (1977), 1798–1799.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY
BATON ROUGE, LA 70817

E-mail address: sbaldrid@math.lsu.edu

MATHEMATICS DEPARTMENT, INDIANA UNIVERSITY
BLOOMINGTON, IN 47405

E-mail address: pkirk@indiana.edu